

DEPARTMENT OF PHYSICS  
INDIAN INSTITUTE OF TECHNOLOGY, MADRAS

PH1020 Physics II

Problem Set 9

19.3.2014

1. An infinitely long cylinder of radius  $a$  has its axis along the  $z$ -axis. Its magnetization is given in cylindrical polar coordinates by  $\mathbf{M} = M_0 (\rho/a)^2 \hat{e}_\phi$ , where  $M_0$  is a constant. Find  $\mathbf{J}_b$  and  $\mathbf{K}_b$  as well as  $\mathbf{B}$  and  $\mathbf{H}$  both inside and outside the cylinder.
2. Consider a toroid in which a wedge-shaped region of small angle  $\psi$  is absent, as shown in the figure. A steady current  $I$  flows in it. The inner radius of the toroid is  $R$ , and the total number of turns in it is  $N$ . Assume that the magnetic field  $\mathbf{B}$  in the air gap is still along  $\hat{e}_\phi$ . Find  $\mathbf{H}$  in the toroid given that the core of the toroid is a LIH magnetic material with magnetic susceptibility  $\chi_m$ .




Figure 1: Top view of the toroid

3. An infinite planar magnetic sheet of thickness  $d$  having a nonuniform permeability given by  $\mu(z) = \mu_0 [1 + (z/d)]^2$  occupies the region  $0 \leq z \leq d$ . There is vacuum on either side of the sheet. A magnetic field  $\mathbf{B} = B_0 \hat{e}_y$  (where  $B_0$  is a constant) is applied in the entire space. The sheet has no free current on it. Find the magnetization surface current densities at  $z = 0$  and  $z = d$ , and also the magnetization volume current density as a function of  $z$ .