

National Programme on Technology Enhanced Learning\*

<http://nptel.iitm.ac.in/courses/115106058/>

Physics – Classical Field Theory

Problem Set 6

<http://sgovindarajan.wikidot.com/cftontheweb>

(To be solved after viewing lecture10 and before lecture 15)

Consider the following Lagrangian density in 1+1 dimensions

$$\mathcal{L} = \frac{1}{2} \partial_\mu \phi \partial^\mu \phi - \frac{\lambda}{2} (\phi^2 - a^2)^2 \quad .$$

1. Obtain the Euler-Lagrange equation corresponding to the Lagrangian density. Verify that the kink soliton given by

$$\phi(x) = a \tanh \mu x \quad ,$$

is a solution of this equation for a particular value of  $\mu$ .

2. Obtain the stress tensor corresponding to the above Lagrangian density.
3. Plot the variation of the Hamiltonian density in space for the kink soliton. In the same plot, plot  $\phi(x)$  for the kink soliton.
4. For the kink soliton, calculate  $E = \int dx \mathcal{H}$  and  $P = \int dx T^{01}$ .
5. By boosting the solution corresponding to the kink soliton, we obtain a new solution of the Euler-Lagrange equation. Verify that this is true. For this solution, now calculate  $E$  and  $P$ . Calculate  $(E^2 - P^2)$  and compare with the unboosted solution.